Recent Surface Changes on Mercury?

- Due to its small size, Mercury was thought to be geologically 'dead' today, except for ongoing impact cratering
- Close-up images from NASA's MESSENGER spacecraft show that some impact craters contain bright shallow depressions, which scientists have named 'hollows'
- The hollows must have formed recently, since they have accumulated few impacts
- Such features have never been seen elsewhere, and the process responsible for their formation remains a mystery

Discoveries in Planetary Science



MESSENGER enhanced-color image of a portion of the Raditladi impact basinon Mercury. The bright, bluish areas are composed of irregular pitlike depressions (hollows). The pits may be actively forming today.

http://dps.aas.org/education/dpsdisc/

A More "Volatile" World Than Expected

- The hollows are found mostly within impact craters, in rocks likely brought up to the surface by an impact event
- Scientists think that these rocks contain abundant *volatile* elements (material that vaporizes easily). *MESSENGER* has measured more volatile elements on Mercury than the Moon.
- High surface temperatures and intense bombardment by the solar wind and micrometeoroids may slowly allow the volatiles to escape the rocks as gas, leading to collapse of the remaining rock



Small hollows on the Sun-facing wall of a 15-km diameter crater. Exposure to the harsh space environment at Mercury is probably responsible for formation of the hollows.

Discoveries in Planetary Science

http://dps.aas.org/education/dpsdisc/

Mercury was expected to form with

- File of the second s
- But this and other MESSENGER results indicate that Mercury has a surprisingly large abundance of volatiles. How did they get there?
- These findings suggest that our ideas about the formation of the planets and the early Solar System need to be revised

Discoveries in Planetary Science



Painting by William K. Hartmann depicting the early Sun and formation of the planets.

MESSENGER results imply that materials originating at different distances from the Sun were mixed extensively to form the planets.

http://dps.aas.org/education/dpsdisc/

For more information...

Press

- New Scientist Sept. 29, 2011 "Bright 'hollows' on Mercury are unique in solar system" http://www.newscientist.com/article/dn20985-bright-hollows-on-mercury-are-unique-to-solar-system.html
- BBC News Sept. 30, 2011 "Hollows' mark Mercury's surface"
 <u>http://www.bbc.co.uk/news/science-environment-15113388</u>

Images

- Slide 1 image courtesy NASA / APL / Carnegie Institute http://photojournal.jpl.nasa.gov/catalog/PIA14856
- Slide 2 image from the source article
- Slide 3 image courtesy William K. Hartmann http://www.psi.edu/~hartmann/catalog/catalog1.html

Source Articles (on-campus login may be required to access journals)

 D.T. Blewett et al., 'Hollows on Mercury: MESSENGER Evidence for Geologically Recent Volatile-Related Activity', Science, 333 (no. 6051), p. 1856-1859, DOI: 10.1126/science.1211681, 2011. http://www.sciencemag.org/content/333/6051/1856.jull/science3328466/-the4-tae5-8004-39ec247764e9

Prepared for the Division for Planetary Sciences of the American Astronomical Society by David Blewett, D. Brain, N. Schneider <u>dpsdisc@aas.org</u> - <u>http://dps.aas.org/education/dpsdisc/</u> - Released November 15, 2012

Discoveries in Planetary Science

http://dps.aas.org/education/dpsdisc/